

*BACILLUS CEREUS*

INTERACTION OF A FREE-LIVING SOIL NEMATODE, *CANENORHABDITIS ELEGANS*,  
WITH SURROGATES OF FOODBORNE PATHOGENIC BACTERIA

(G. L. Anderson, K. N. Caldwell, L. R. Beuchat, and P. L. Williams)

The agricultural impacts of plant and animal parasitic nematodes have long been recognized and, by virtue of their effects on fruit and vegetable production, have been extensively studied. Comparatively little is known regarding the impact of free-living microbiovorous nematodes on produce production and safety, although they are the most abundant and wide spread soil mesofauna. The association of free-living nematodes and various genera of bacteria has been studied. While it is recognized that free-living nematodes avoid certain bacteria, it is clear that they do not uniformly avoid foodborne pathogens. Two human enteric pathogens, *Salmonella* and *Shigella*, are reportedly ingested and defecated by free-living saprozoic nematodes and *Salmonella* Typhimurium is known to infect the free-living soil nematode *Caenorhabditis elegans*. From these reports, it appears that free-living nematodes may be important as vectors of pathogenic bacteria, including some forms capable of causing human disease.

Soil is a source of microbial contamination of fruit and vegetables, as evidenced by the isolation of soil-residing pathogenic bacteria from produce. In a survey of vegetables for the presence of amoebae and *Salmonella*, nematode eggs and larvae have been recovered using a naccional-ether method. The recovery of nematodes from uncooked vegetables indicates that agronomic conditions and marketing practices may be conducive to the survival of nematodes on fresh produce. This also indicates that if free-living nematodes are present on raw produce, they may serve as vehicles for contamination with pathogenic bacteria, either by contact with their surface or via eggs or voided material from their gastrointestinal tract.

We undertook a study to evaluate the interaction of *C. elegans* with bacterial surrogates for foodborne pathogens occasionally occurring or persisting in soil. Nematode/bacterial interactions were characterized to determine the propensity of young adult worms to be attracted to bacterial colonies, to compare the feeding and development of young adult worms cultured on this diverse group of bacteria, and to examine the dispersal of bacteria by *C. elegans* following feeding on monoxenic cultures. We evaluated the association between a free-living soil nematode, *C. elegans*, with *Escherichia coli*, an avirulent strain of *S. Typhimurium*, *Listeria welshimeri*, and *Bacillus cereus*. On an agar medium, young adult worms quickly moved toward colonies of all four bacteria; over 90% of 3-day adults entered colonies within 16 min after inoculation. After 48 h, worms moved in and out of colonies of *L. welshimeri* and *B. cereus*, but remained associated with *E. coli* and *S. Typhimurium* colonies for at least 96 h. Young adult worms fed on cells of the four bacteria suspended in K medium. Worms survived and reproduced using nutrients derived from all test bacteria, as evidenced by eggs laid by second generation worms after culturing for 96 h. Development was slightly slower in worms fed on Gram-positive bacteria compared to Gram-negative bacteria. Worms fed for 24 h on bacterial lawns formed on tryptic soy agar dispersed bacteria over a 3-h period when transferred to a bacteria-free agar surface. Results suggest that *C. elegans* and, perhaps, other free-living nematodes are potential vectors for both Gram-positive and Gram-negative bacteria, including foodborne pathogens in soil.